



Clinical Audit

# An audit on pediatric cholesteatoma

S. Lakshmi Priya

ENT, Kauvery Hospital, Trichy, Cantonment, Trichy

\*Correspondence

**Abstract:** Cholesteatoma is a complex condition characterized by the abnormal presence of keratinizing squamous epithelium in the middle ear or mastoid, accompanied by subepithelial connective tissue and accumulating keratin debris. Despite its misleading name, cholesteatoma contains neither cholesterol crystals nor tumor tissue. This condition is particularly aggressive in children and, if left untreated, can lead to bone erosion and serious complications. The pathophysiology of bone resorption in cholesteatoma involves multiple mechanisms, including enzymatic-induced necrosis, cytokine-mediated bone remodeling, prostaglandin-induced changes, and potential bacterial involvement. This study aims to describe our methodology and outcomes in treating pediatric patients with cholesteatoma, addressing the unique challenges posed by the condition in this age group. By examining our approach and results, we hope to contribute to the development of more effective management strategies for pediatric cholesteatoma, ultimately improving patient outcomes and reducing the risk of complications.

**Keywords:** Ear disease; otitis media; chronic otitis media; cochlear implants; hearing loss

## 1. Introduction

The ear is a complex sensory organ composed of three main parts: the external ear, middle ear, and internal ear. The external ear consists of the visible portion, known as the pinna, and the ear canal. The middle ear, a small air-filled cavity, houses three tiny bones called ossicles: the malleus (hammer), incus (anvil), and stapes (stirrup). These ossicles form a chain that transmits sound vibrations from the eardrum to the inner ear. The middle ear is further divided into three regions: the epitympanum (upper portion), mesotympanum (middle portion), and hypotympanum (lower portion). This anatomical arrangement allows for efficient sound conduction and amplification. The internal ear, also known as the inner ear, contains the cochlea and semicircular canals, which are responsible for converting sound waves into electrical signals and maintaining balance, respectively.

Facial nerve leaves the brainstem at pontomedullary junction; it travels through the posterior cranial fossa and enters the IAM. At the fundus of the IAM, it takes a turn posteriorly forming a genu, from the genu the nerve passes backward above the oval window and below the SCC till the level of pyramidal eminence (2nd genu). The nerve continues vertically downward along the posterior wall of tympanic cavity and leaves the temporal bone through the stylomastoid foramen

### What is cholesteatoma?

Cholesteatoma

The presence of keratinized squamous epithelium in middle ear or mastoid is characteristic of cholesteatoma – “Skin in the wrong place”

The term ‘cholesteatoma’ is a misnomer; it contains neither cholesterol crystals nor a tumor.

**Citation:** Lakshmi Priya S. An audit on pediatric cholesteatoma. *Kauverian Med J.*, 2024;1(12):1-12.

Academic Editor: Dr. Venkita S. Suresh

ISSN: 2584-1572 (Online)




**Copyright:** © 2024 by the authors. Submitted for possible open access publication under the terms and conditions.

### Definition of Cholesteatoma

Cholesteatoma is a mass formed by keratinizing squamous epithelium in the middle ear and/or mastoid, subepithelial connective tissue and by progressive accumulation of keratin debris with/without surrounding inflammatory reaction.

It is aggressive in children, if left untreated, can cause bone erosion which in turn can lead on to serious complications.



### Bone resorption in Cholesteatoma

- 1) Enzymatic-induced necrosis and resorption - Collagenase, Acid phosphatase, proteolytic enzymes
- 2) Cytokines induced bone remodelling – IL-1a and 1b, TNF-alpha, TNF-beta
- 3) Prostaglandin induced bone remodelling - PG E2
- 4) Pyogenic osteitis theory – Pseudomonas aeruginosa, Streptococcus sp, proteus, E.coli
- 5) Pressure necrosis theory – unlikely

### Objectives

To describe our methodology and outcome in treating pediatric patients with Cholesteatoma.

### 2. Materials and Methods

- 1) A retrospective study was carried out from July 2023–June 2024.
- 2) The study includes 10 patients, of whom
  - 4 were boys
  - 6 were girls
- 3) Between the age of 7 and 16 years.


The evaluated parameters

- 1) Clinical symptoms
- 2) Site of infection
- 3) Extension of the disease

- 4) Ossicular chain involvement
- 5) Facial nerve integrity
- 6) Imaging
- 7) Treatment modality
- 8) Outcome
- 9) Complications

### Symptoms and Signs

#### FACIAL ASSESSMENT



- 1) All patients invariably presented with Foul smelling ear discharge and Hard of hearing.
- 2) 50% of patients had blood stained ear discharge.
- 3) 20% of patients had facial asymmetry.

### Preoperative Assessment of Facial Nerve



- a. House-Brackmann grade 4
- b. He also had Giddiness, Vomiting, Headache



- (1) - Complete left eye closure only with maximum effort
  - (2) - Asymmetry of mouth with maximum effort
  - (3) - Inability to blow the cheek
- House-Brackmann grade 3

**Otoendoscopy**




Normal TM




Retracted TM with eroded incus

■ Attic retraction ■ PSQ retraction




■ Ossicular erosion    ■ Intact ossicle



**CT Temporal Bone – Axial Plane**

Soft tissue mass extending deep to involve the semicircular canals, encompassing the labyrinthine segment of fallopian canal and causing erosion of internal auditory meatus



**CT Temporal Bone – Coronal Plane**

Erosion of Lateral epitympanic wall (the scutum). Erosion of ossicles




Sagging of Posterior wall of external auditory canal Soft tissue mass in middle ear



### CT Temporal Bone – Coronal Plane

Extensive bone destruction of the mastoid mimicking the appearance of mastoidectomy–Automastoidectomy



### Diagnosis


Diagnosis	No. of. Cases
Chronic otitis media with cholesteatoma, active, squamosal disease with facial nerve compression	2
Chronic otitis media with cholesteatoma, active, squamosal disease with intact facial nerve	8

### Management


Surgery	No. of. Cases
Canal wall down mastoidectomy, tympanoplasty with facial nerve decompression	2
Canal wall down mastoidectomy with tympanoplasty	8

### Intraoperative Findings


Cholesteatoma Site of Involvement



**Ossicular status**



**Fallopian Canal and Semicircular Canal Status**



**Left ear CWD Mastoidectomy cavity of a 7-year-old girl**

Handle of malleus with head being nipped off  
 Dome of horizontal semicircular canal  
 Chorda tympani nerve  
 Head of stapes




**Ossicular Chain Reconstruction**

Sculptured head of malleus placed over head of stapes



Graft materials	No.of. Cases	Placed over	Tympanoplasty type
Harvested conchal cartilage	2	Head of stapes	III
Septal cartilage	1	Head of stapes	III
Sculptured Malleus	1	Head of stapes	III
Sculptured Incus	1	Head of stapes	III
Sculptured Incus	2	Footplate of stapes	IV
PORP	1	Head of stapes	III
TORP	1	Footplate of stapes	IV




**Left ear CWD Mastoidectomy of a 9-year-old boy**


Cholesteatoma flakes in supratubal recess



Disease clearance from sinus tympani




Granulation tissue being dissected from the head of stapes






Cholesteatoma flakes filling the mastoid tip cells




Mastoid cavity after disease clearance



### Right ear CWD Mastoidectomy of a 16-year-old girl

Dome of horizontal semicircular canal eroded in the superior aspect  
Facial nerve being exposed from 1st genu till chorda tympani branching  
Cholesteatoma matrix in superior semicircular canal



### Pre OP – HB Grade 4



**Post OP – Normal**




**Pre OP – HB Grade 3**



**Post OP – Normal**



**Otoendoscopy of Postoperative Mastoid Cavity**



**Comparison**

<b>Variables</b>	<b>Our study</b>	<b>Acta Ital</b>
Number of patients	10 in 1 year	36 in 6 years
Age group	7–16 years	6–14 years
<b>Surgery</b>		
Canal wall down mastoid-ectomy	100%	39.39%
Canal wall intact mastoid-ectomy	0	60.60%
<b>Complications</b>		
Residual disease	Nil	18.18%
Recurrence	Nil (On Avg 5 months follow-up)	6.06%
<b>Hearing</b>		
Subjective test	Hearing improvement present	Hearing improvement present
Objective test (Audiometry)	Not done	ABG between 0 and 20 db in 54%

**Surgery**

<b>Primary role of surgery</b>	<b>Secondary role of surgery</b>
Remove all original cholesteatoma squamous epithelial matrix Prevent further erosion and complications Give a dry, watertight ear Give a ear that will be self-cleaning Prevent the occurrence of recurrent cholesteatoma.	Improve hearing (aim is to restore the best possible hearing in an ear that will never be normal).

<b>What to be cautious in pediatric Cholesteatoma?</b>	<b>Why/how?</b>
More extensive and destructive in children than adults Increases risk of complications Impact on development Difficulty in diagnosis Surgical challenges High recurrence rates Impact on hearing and balance Potential for congenital associations	Hence early intervention is important to prevent further damage More prone to facial paralysis, Meningitis, Brain abscess if left untreated, it affects speech, language and cognitive development Due to nonspecific symptoms Surgery can be more complex in children due to extensive spread in developing temporal bone More likely to recur due to poor Eustachian tube function. Hence long term follow-up will be necessary As it cause erosion of ossicles and labyrinth, patient develops hard of hearing and giddiness respectively Down’s syndrome, Craniofacial syndrome, di-George syndrome

---

Emotional and psychological impact

Children and families may experience emotional distress and concerns about hearing

---

### **3. Conclusion**

Management of pediatric cholesteatoma requires a highly individualized approach that takes into account anatomic, clinical and social factors to determine the most successful treatment paradigm.

Prompt evaluation, management, and early intervention are very important to avoid life-threatening events.